



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/00	A2	(11) International Publication Number: WO 98/25958 (43) International Publication Date: 18 June 1998 (18.06.98)
(21) International Application Number: PCT/US97/22766 (22) International Filing Date: 12 December 1997 (12.12.97) (30) Priority Data: 60/032,846 13 December 1996 (13.12.96) US (71) Applicant: SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US). (72) Inventors: GORMAN, Daniel, M.; 6371 Central Avenue, Newark, CA 94560 (US). MATTSON, Jeanine, D.; 559 Alvarado Street, San Francisco, CA 94114 (US). (74) Agents: THAMPOE, Immac, J. et al.; Schering-Plough Corporation, Patent Dept., K-6-1, 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CZ, EE, GE, GW, HU, ID, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: MAMMALIAN CELL SURFACE ANTIGENS; RELATED REAGENTS (57) Abstract Purified genes encoding a T cell surface antigen from a mammal, reagents related thereto including purified proteins, specific antibodies, and nucleic acids encoding this antigen are provided. Methods of using said reagents and diagnostic kits are also provided.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NI	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China						

MAMMALIAN CELL SURFACE ANTIGENS; RELATED REAGENTS

5

FIELD OF THE INVENTION

The present invention pertains to compositions related to proteins which function in controlling activation and expansion of mammalian cells, e.g., cells of a mammalian immune system. In particular, it provides purified genes, proteins, antibodies, and related reagents useful, e.g., to regulate activation, development, differentiation, and function of various cell types, including hematopoietic cells.

BACKGROUND OF THE INVENTION

The activation of resting T cells is critical to most immune responses and allows these cells to exert their regulatory or effector capabilities. See Paul (ed; 1993) Fundamental Immunology 3d ed., Raven Press, N.Y. Increased adhesion between T cells and antigen presenting cells (APC) or other forms of primary stimuli, e.g., immobilized monoclonal antibodies (mAb), can potentiate the T-cell receptor signals. T-cell activation and T cell expansion depends upon engagement of the T-cell receptor (TCR) and co-stimulatory signals provided by accessory cells. See, e.g., Jenkins and Johnson (1993) Curr. Opin. Immunol. 5:361-367; Bierer and Hahn (1993) Semin. Immunol. 5:249-261; June, et al. (1990) Immunol. Today 11:211-216; and Jenkins (1994) Immunity 1:443-446. A major, and well-studied, co-stimulatory interaction for T cells involves either CD28 or CTLA-4 on T cells with either B7 or B70 (Jenkins (1994) Immunity 1:443-446). Recent studies on CD28 deficient mice (Shahinian, et al. (1993) Science 261:609-612; Green, et al. (1994) Immunity

2

transgenic mice (Ronchese, et al. (1994) J. Exp. Med. 179:809-817) have revealed deficiencies in some T-cell responses though these mice have normal primary immune responses and normal CTL responses to lymphocytic choriomeningitis virus and vesicular stomatitis virus. As a result, both these studies conclude that other co-stimulatory molecules must be supporting T-cell function. However, identification of these molecules which mediate distinct costimulatory signals has been difficult.

10 Tumor Necrosis Factor (TNF) is the prototypic member of an emerging family of cytokines that function as prominent mediators of immune regulation and the inflammatory response. These ligands are typically type II membrane proteins, with homology at the carboxy terminus. A proteolytic processed soluble protein often is produced. See, e.g., Smith, et al. (1994) Cell 76-959-962; Armitage (1994) Current Opinion in Immunology 6:407-413; Gruss and Dower (1995) Blood 85:3378-3404; Wiley, et al. (1995) Immunity 3:673-682; and Baker and Reddy (1996) Oncogene 12:1-9. Crucial roles for these family members are evidenced by a number of studies, and they are implicated in regulation of apoptosis, peripheral tolerance, Ig maturation and isotype switching, and general B cell and T cell functions. See, e.g., Thomson (ed. 1994) The cytokine Handbook Academic Press, San Diego, CA. These imply fundamental roles in immune and developmental networks.

30 The inability to modulate activation signals prevents control of inappropriate developmental or physiological responses in the immune system. The present invention provides at least one alternative costimulatory molecule, agonists and antagonists of which will be useful in modulating a plethora of immune responses.

3

SUMMARY OF THE INVENTION

The present invention is based, in part, upon the discovery of an antigen which exhibits sequence homology to proteins which act as inducers of apoptosis. In particular, it provides a gene encoding a 316 amino acid protein, designated 499E9, which is expressed on a highly polarized Th1 T cell. Engagement of 499E9 may modulate antigen-specific proliferation and cytokine production by effector cells. 499E9 is a novel cell surface molecule which, when engaged, may either potentiate immune cell expansion or apoptosis. The mouse embodiment is described, enabling mammalian genes, proteins, antibodies, and uses thereof. Functional equivalents exhibiting significant sequence homology are available from other mammalian, e.g., human, and non-mammalian species. Moreover, the receptor of 499E9 can function as its binding partner to stimulate other cells expressing the receptor.

More particularly, the present invention provides a composition of matter selected from: a substantially pure or recombinant 499E9 protein or peptide exhibiting at least about 85% sequence identity over a length of at least about 12 amino acids to SEQ ID NO: 2; a natural sequence 499E9 of SEQ ID NO: 2; or a fusion protein comprising 499E9 sequence. Certain embodiments include a substantially pure or isolated protein comprising a segment exhibiting sequence identity to a corresponding portion of a 499E9, wherein: the homology is at least about 90% identity and the portion is at least about 9 amino acids; the homology is at least about 80% identity and the portion is at least about 17 amino acids; or the homology is at least about 70% identity and the portion is at least about 25 amino acids. Other embodiments include a composition of matter, wherein the: 499E9 comprises a mature sequence of Table 1; or protein or peptide: is from a warm blooded animal selected from a

4

polypeptide segment of SEQ ID NO: 2; exhibits a plurality of portions exhibiting the identity; is a natural allelic variant of 499E9; has a length at least about 30 amino acids; exhibits at least two non-overlapping epitopes

5 which are specific for a mammalian 499E9; exhibits a sequence identity at least about 90% over a length of at least about 20 amino acids to a rodent 499E9; exhibits at least two non-overlapping epitopes which are specific for a rodent 499E9; exhibits a sequence identity at least

10 about 90% over a length of at least about 20 amino acids to a rodent 499E9; is glycosylated; is a synthetic polypeptide; is attached to a solid substrate; is conjugated to another chemical moiety; is a 5-fold or less substitution from natural sequence; or is a deletion

15 or insertion variant from a natural sequence. Also provided are various compositions, e.g., comprising: a sterile 499E9 protein or peptide; or the 499E9 protein or peptide and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or

20 formulated for oral, rectal, nasal, topical, or parenteral administration. Fusion proteins are provided, e.g., comprising: mature protein sequence of Table 1; a detection or purification tag, including a FLAG, His6, or Ig sequence; or sequence of another TNF-ligand protein.

25 Kit embodiments are provided, e.g., comprising such a protein or polypeptide, and: a compartment comprising the protein or polypeptide; and/or instructions for use or disposal of reagents in the kit.

Antibody, or binding compound embodiments include

30 those comprising an antigen binding portion from an antibody, which specifically binds to a natural 499E9 protein, wherein: the protein is a rodent protein; the binding compound is an Fv, Fab, or Fab2 fragment; the binding compound is conjugated to another chemical

35 moiety; or the antibody: is raised against a peptide sequence of a mature polypeptide comprising sequence of

- purified 499E9; is immunoselected; is a polyclonal antibody; binds to a denatured 499E9; exhibits a Kd to antigen of at least 30 μ M; is attached to a solid substrate, including a bead or plastic membrane; is in a sterile composition; or is detectably labeled, including a radioactive or fluorescent label. Other embodiments include a kit comprising the binding compound, and: a compartment comprising the binding compound; and/or instructions for use or disposal of reagents in the kit.
- 10 Other forms include, e.g., a composition comprising: a sterile binding compound; or the binding compound and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for oral, rectal, nasal, topical, or parenteral
- 15 administration. Such also allow methods of purifying a 499E9 protein or peptide from other materials in a mixture comprising contacting the mixture to such an antibody, and separating bound 499E9 from other materials.
- 20 Nucleic acid embodiments include an isolated or recombinant nucleic acid encoding a protein or peptide or fusion protein, wherein: the 499E9 protein is from a mammal, including a rodent; or the nucleic acid: encodes an antigenic peptide sequence of Table 1; encodes a
- 25 plurality of antigenic peptide sequences of Table 1; exhibits at least about 80% identity to a natural cDNA encoding the segment; is an expression vector; further comprises an origin of replication; is from a natural source; comprises a detectable label; comprises synthetic
- 30 nucleotide sequence; is less than 6 kb, preferably less than 3 kb; is from a mammal, including a rodent; comprises a natural full length coding sequence; is a hybridization probe for a gene encoding the 499E9 protein; or is a PCR primer, PCR product, or mutagenesis
- 35 primer. A cell or tissue comprising such a recombinant nucleic acid is also embraced within the invention, e.g.,

- cell; a bacterial cell; a yeast cell; an insect cell; a mammalian cell; a mouse cell; a rodent cell; or a human cell. Kit forms include those comprising the nucleic acid, and: a compartment comprising the nucleic acid; a
- 5 compartment further comprising a 499E9 protein or polypeptide; and/or instructions for use or disposal of reagents in the kit. Other nucleic acid embodiments include those which: hybridize under wash conditions of: 30° C and less than 2M salt, 45° C and/or 500 mM salt, or
- 10 at 55° C and/or 150 mM salt to SEQ ID NO: 1; or exhibit at least about 85% identity over a stretch of at least about 30 nucleotides, at least 90% and/or the stretch is at least 55 nucleotides, or at least 95% and/or the stretch is at least 75 nucleotides to a rodent 499E9.
- 15 The invention further provides methods of modulating physiology or development of a cell or tissue culture cells comprising introducing into the cell an agonist or antagonist of a 499E9. Other methods include modulating the physiology of a cell comprising contacting the cell
- 20 with: a substantially pure 499E9 or fragment; an antibody or binding partner which specifically binds a 499E9; or a nucleic acid encoding a 499E9 or peptide. Preferably, the cell is a T cell and the modulating of physiology is: apoptosis of the T cell; or activation of the T cell.
- 25 The invention further provides a method of treating a patient having an abnormal immune response by administering an effective dose of an antibody or binding partner specific for 499E9; a 499E9 protein or polypeptide; or a nucleic acid encoding a 499E9 peptide.
- 30 The abnormal immune response is characterized by a T cell immune deficiency; chronic inflammation; or tissue rejection.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

All references cited herein are incorporated herein by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

I. General

The present invention provides amino acid sequences and DNA sequences encoding various mammalian proteins which are antigens found in many T cell subtypes, e.g., Th1, Th2, polarized Th1 cells, and polarized Th2 cells. Among these proteins are antigens which modulate, e.g., induce or prevent proliferation or differentiation of interacting cells, among other physiological effects. The full length antigens, and fragments, or antagonists will be useful in physiological modulation of cells expressing counter receptors for the antigen. The proteins will also be useful as antigens, e.g., immunogens, for raising antibodies to various epitopes on the protein, both linear and conformational epitopes. The molecule may be useful in defining functional T cell or NK cell subsets.

A cDNA encoding 499E9 was isolated from a polarized Th1 cell cDNA library, see Openshaw, et al. (1995) J. Exp. Med. 182:1357-1367. The 499E9 cDNA contains a stretch of about 2191 bp in length and contained one large open reading frame encoding a type II transmembrane protein. Transcript analysis has identified multiple transcripts with the most prevalent being 2.1 to 2.3 kb. Structural features include an intracellular domain sequence of about 52 amino acids, an extracellular region of about 246 amino acids, and a hydrophobic presumptive membrane spanning portion of about 20 amino acids. See Table 1 and SEQ. ID. NO: 2. 499E9 exhibits structural motifs characteristic of a member of the TNF ligand

ligand, TNF, NGF, and FAS. Table 1 illustrates the nucleic acid and predicted amino acid sequences for mouse 499E9.

5

Table 1: Mouse 499E9 nucleotide and predicted amino-acid sequence. Predicted intracellular domain sequence runs about from met1 to met49; residues 8 and 11 are potential tyrosine phosphorylation sites; a transmembrane sequence probably runs about from phe50 to leu69; and the extracellular domain probably runs about from tyr70 to asp316. See SEQ ID NO: 1 and 2.

10

GCCAGGACCT CTGTGAACCG GTCGGGGCGG GGGCCGCCTG GCCGGGAGTC TGCTCGGCGG
60

15

TGGGTGGCCG AGGAACGGAG AGAACGATCG CGGAGCAGGG CGCCCGAACT CCGGGCGCCG
120

20

CGCC ATG CGC CGG GCC AGC CGA GAC TAC GGC AAG TAC CTG CGC AGC TCG
169
Met Arg Arg Ala Ser Arg Asp Tyr Gly Lys Tyr Leu Arg Ser Ser
1 5 10 15

25

GAG GAG ATG GGC AGC GGC CCC GGC GTC CCA CAC GAG GGT CCG CTG CAC
217
Glu Glu Met Gly Ser Gly Pro Gly Val Pro His Glu Gly Pro Leu His
20 25 30

30

CCC GCG CCT TCT GCA CCG GCT CCG GCG CCG CCA CCC GCC GCC TCC CGC
265
Pro Ala Pro Ser Ala Pro Ala Pro Ala Pro Pro Pro Ala Ala Ser Arg
35 40 45

35

TCC ATG TTC CTG GCC CTC CTG GGG CTG GGA CTG GGC CAG GTG GTC TGC
313
Ser Met Phe Leu Ala Leu Leu Gly Leu Gly Leu Gly Gln Val Val Cys
50 55 60

40

AGC ATC GCT CTG TTC CTG TAC TTT CGA GCG CAG ATG GAT CCT AAC AGA
361
Ser Ile Ala Leu Phe Leu Tyr Phe Arg Ala Gln Met Asp Pro Asn Arg
65 70 75

45

ATA TCA GAA GAC AGC ACT CAC TGC TTT TAT AGA ATC CTG AGA CTC CAT
409
Ile Ser Glu Asp Ser Thr His Cys Phe Tyr Arg Ile Leu Arg Leu His
80 85 90 95

50

GAA AAC GCA GGT TTG CAG GAC TCG ACT CTG GAG AGT GAA GAC ACA CTA
457
Glu Asn Ala Gly Leu Gln Asp Ser Thr Leu Glu Ser Glu Asp Thr Leu
100 105 110

9

CCT GAC TCC TGC AGG AGG ATG AAA CAA GCC TTT CAG GGG GCC GTG CAG
 505
 5 Pro Asp Ser Cys Arg Arg Met Lys Gln Ala Phe Gln Gly Ala Val Gln
 115 120 125

AAG GAA CTG CAA CAC ATT GTG GGG CCA CAG CGC TTC TCA GGA GCT CCA
 553
 10 Lys Glu Leu Gln His Ile Val Gly Pro Gln Arg Phe Ser Gly Ala Pro
 130 135 140

GCT ATG ATG GAA GGC TCA TGG TTG GAT GTG GCC CAG CGA GGC AAG CCT
 601
 15 Ala Met Met Glu Gly Ser Trp Leu Asp Val Ala Gln Arg Gly Lys Pro
 145 150 155

GAG GCC CAG CCA TTT GCA CAC CTC ACC ATC AAT GCT GCC AGC ATC CCA
 649
 20 Glu Ala Gln Pro Phe Ala His Leu Thr Ile Asn Ala Ala Ser Ile Pro
 160 165 170 175

TCG GGT TCC CAT AAA GTC ACT CTG TCC TCT TGG TAC CAC GAT CGA GGC
 697
 25 Ser Gly Ser His Lys Val Thr Leu Ser Ser Trp Tyr His Asp Arg Gly
 180 185 190

TGG GCC AAG ATC TCT AAC ATG ACG TTA AGC AAC GGA AAA CTA AGG GTT
 745
 30 Trp Ala Lys Ile Ser Asn Met Thr Leu Ser Asn Gly Lys Leu Arg Val
 195 200 205

AAC CAA GAT GGC TTC TAT TAC CTG TAC GCC AAC ATT TGC TTT CGG CAT
 793
 35 Asn Gln Asp Gly Phe Tyr Tyr Leu Tyr Ala Asn Ile Cys Phe Arg His
 210 215 220

CAT GAA ACA TCG GGA AGC GTA CCT ACA GAC TAT CTT CAG CTG ATG GTG
 841
 40 His Glu Thr Ser Gly Ser Val Pro Thr Asp Tyr Leu Gln Leu Met Val
 225 230 235

TAT GTC GTT AAA ACC AGC ATC AAA ATC CCA AGT TCT CAT AAC CTG ATG
 889
 45 Tyr Val Val Lys Thr Ser Ile Lys Ile Pro Ser Ser His Asn Leu Met
 240 245 250 255

AAA GGA GGG AGC ACG AAA AAC TGG TCG GGC AAT TCT GAA TTC CAC TTT
 937
 50 Lys Gly Gly Ser Thr Lys Asn Trp Ser Gly Asn Ser Glu Phe His Phe
 260 265 270

TAT TCC ATA AAT GTT GGG GGA TTT TTC AAG CTC CGA GCT GGT GAA GAA
 985
 Thr Ser Thr Asn Val Gly Gly Phe Phe Lys Leu Arg Ala Gly Gly Gly

ATT AGC ATT CAG GTG TCC AAC CCT TCC CTG CTG GAT CCG GAT CAA GAT
 1033
 Ile Ser Ile Gln Val Ser Asn Pro Ser Leu Leu Asp Pro Asp Gln Asp
 5 290 295 300

GCG ACG TAC TTT GGG GCT TTC AAA GTT CAG GAC ATA GAC TGAGACTCAT
 1082
 Ala Thr Tyr Phe Gly Ala Phe Lys Val Gln Asp Ile Asp
 10 305 310 315

TTCGTGGAAC ATTAGCATGG ATGTCCTAGA TGTTTGGAAC CTTCTTAAAA AATGGATGAT
 1142

GTCTATACAT GTGTAAGACT ACTAAGAGAC ATGGCCCACG GTGTATGAAA CTCACAGCCC
 1202

TCTCTCTTGA GCCTGTACAG GTTGTGTATA TGTAAGTCC ATAGGTGATG TTAGATTTCAT
 1262

GGTGATTACA CAACGGTTTTC ACAATTTTGT AATGATTTC TAAGAATTGA ACCAGATTTC
 1322

GAGAGGTATT CCGATGCTTA TGAAAACTT ACACGTGAGC TATGGAAGGG GGTCACAGTC
 1382

TCTGGGTCTA ACCCCTGGAC ATGTGCCACT GAGAACCCTG AAATTAAGAA GATGCCATGT
 1442

CATTGCAAAG AAATGATAGT GTGAAGGGTT AAGTTCTTTT GAATTGTTAC ATTGCGCTGG
 1502

GACCTGCAAA TAAGTTCTTT TTTTCTAATG AGGAGAGAAA AATATATGTA TTTTATATA
 1562

ATGTCTAAAG TTATATTTCA GGTGTAATGT TTTCTGTGCA AAGTTTGTGA AATTATATTT
 1622

GTGCTATAGT ATTTGATTCA AAATATTTAA AAATGTCTCA CTGTTGACAT ATTTAATGTT
 1682

TTAAATGTAC AGATGTATTT AACTGGTGCA CTTTGTAAAT CCCCTGAAGG TACTCGTAGC
 1742

TAAGGGGGCA GAATACTGTT TCTGGTGACC ACATGTAGTT TATTTCTTTA TTCTTTTAA
 1802

CTTAATAGAG TCTTCAGACT TGTCAAAACAT ATGCAAGCAA AATAAATAAA TAAAAATAAA
 1862

ATGAATATCT TGAATAATAA GTAGGATGTT GGTCACCAGG TGCCTTTCAA ATTTAGAAGC
 1922

TCAGGAGTAT TTATGCAATT ATTGAACAGG TGTCTTTTTT TACAAGAGCT ACAAATTGTA
2042

5 AATTTTGTTT CTTTTTTTTC CCATAGAAAA TGTACTATAG TTTATCAGCC AAAAAACAAT
2102

CCACTTTTTTA ATTTAGTGAA AGTTATTTTA TTATACTGTA CAATAAAAGC ATTGTTTCTG
2162

10 AATGGCATT TTTGGTACTT AAAAATGGC
2191

TNF ligand family members have a conserved leucine
15 residue corresponding to residue 205; a conserved glycine
residue corresponding to residue 211; a conserved
tyrosine residue corresponding to residue 216; a
conserved glycine residue corresponding to residue 277; a
conserved leucine residue corresponding to residue 282; a
20 conserved phenylalanine residue corresponding to 307; and
a conserved glycine residue corresponding to residue 308.
The TNF ligand domain seems to run about from 205 (leu)
to 316 (asp). Glycosylation sites may be at 197 and 262.
This clone exhibits closes homology to a mouse TRAIL,
25 which is implicated in induction of apoptosis. Related
family members include ligands for CD40 and FAS, and
lymphotoxin beta, tumor necrosis factor, etc.

By cDNA Southern analysis, it is clear that 499E9 is
expressed in many T cells, including Th1, Th2, 3 week
30 polarized Th1 or Th2 cells, pre T cells, and in Rag
knock-out thymus cDNA libraries. Some weak signal from
dendritic cells may have been detected. Cells expressing
499E9 typically contain a main transcript of about 2.1 to
2.3 kb, but also containing other transcripts. Tissue
35 distribution analysis suggests a positive signal in
brain, heart, kidney, liver, lung, spleen, and testis.
Transcripts for 499E9 have not been detected in
fibroblasts (L cells), monocytes (RAW264), naive T cells
(CD4+, MEL14+, Br cells), macrophage cells, Nippo
40 infected lung/liver/spleen, or Rag knock out organs

The structural homology of 499E9 to the TNF ligand family suggests function of this molecule. 499E9, as a T cell surface molecule, likely modulates Ag-specific proliferative responses on effector cells, or induction
5 of apoptosis of those cells. 499E9 agonists, or antagonists, may also act as a co-stimulatory molecule for regulation of T cell mediated cell activation, and may in fact, cause a shift of T helper cell types, e.g., between Th1 and Th2. Thus, 499E9 or antagonists should
10 be useful in the treatment of abnormal immune disorders, e.g., T cell immune deficiencies, chronic inflammation, or tissue rejection.

TNF ligand molecules typically modulate cell proliferation, viability, and differentiation. For
15 example, TNF and FAS can kill cells expressing their respective receptors, including fibroblasts, liver cells, and lymphocytes. Some members of this class of ligands exhibit effects on cellular proliferation of cells expressing their respective receptors, e.g., B cells
20 expressing CD40. These effects on proliferation may also effect subsequent differentiation steps, and may lead, directly or indirectly, to changes in cytokine expression profiles.

The members of the TNF ligand family also exhibit
25 costimulation effects, which may also regulate cellular differentiation or apoptosis. Receptor expressing cells may be protected from activation induced cell death (AICD) or apoptosis. For example, CD40 ligand can have effects on T and B lymphocytes.

30 The embodiment characterized herein is from mouse, but other primate, e.g., human, variants will exist. Additional sequences for proteins in other mammalian species, e.g., primates and rodents, will also be available. See below. The descriptions below are
35 directed, for exemplary purposes, to a mouse 499E9, but are likewise applicable to related embodiments from other

The mouse 499E9 protein is a protein which exhibits structural features characteristic of a cell surface antigen, e.g., a TNF ligand family member. The protein is easily detected on particular cell types, others
5 express lesser amounts. The 499E9 antigen should be present in the identified tissue types and the interaction of the antigen with its binding partner should be important for mediating various aspects of cellular physiology or development, as described.

10

II. Purified 499E9

Mouse 499E9 amino acid sequence is shown in SEQ ID NO: 2. These amino acid sequences, provided amino to carboxy, are important in providing sequence information
15 in the antigen allowing for distinguishing the protein from other proteins and exemplifying numerous variants. Moreover, the peptide sequences allow preparation of peptides to generate antibodies to recognize such segments, and nucleotide sequences allow preparation of
20 oligonucleotide probes, both of which are strategies for detection or isolation, e.g., cloning, of genes or cDNAs encoding such sequences.

As used herein, the term "mouse 499E9" shall encompass, when used in a protein context, a protein
25 having amino acid sequence shown in SEQ ID NO: 2, or a significant fragment of such a protein, or another highly homologous protein derived from mouse. These binding components, e.g., antibodies, typically bind to a 499E9 with high affinity, e.g., at least about 100 nM, usually
30 better than about 30 nM, preferably better than about 10 nM, and more preferably at better than about 3 nM. Homologous proteins would be found in mammalian species other than mouse, e.g., primates or rodents. Non-mammalian species should also possess structurally or
35 functionally related genes and proteins, e.g., birds or amphibians.

14

The term "polypeptide" as used herein includes a significant fragment or segment, and encompasses a stretch of amino acid residues of at least about 8 amino acids, generally at least about 12 amino acids, typically at least about 16 amino acids, preferably at least about 20 amino acids, and, in particularly preferred embodiments, at least about 30 or more amino acids.

The term "binding composition" refers to molecules that bind with specificity to 499E9, e.g., in a cell adhesion pairing type fashion, or an antibody-antigen interaction. It also includes compounds, e.g., proteins, which specifically associate with 499E9, including in a natural physiologically relevant protein-protein interaction, either covalent or non-covalent. The molecule may be a polymer, or chemical reagent. A functional analog may be an antigen with structural modifications, or it may be a molecule which has a molecular shape which interacts with the appropriate binding determinants. The compounds may serve as agonists or antagonists of the binding interaction, see, e.g., Goodman, et al. (eds.) (1990) Goodman & Gilman's: The Pharmacological Bases of Therapeutics (8th ed.), Pergamon Press.

Substantially pure typically means that the protein is free from other contaminating proteins, nucleic acids, or other biologicals derived from the original source organism. Purity may be assayed by standard methods, typically by weight, and will ordinarily be at least about 40% pure, generally at least about 50% pure, often at least about 60% pure, typically at least about 80% pure, preferably at least about 90% pure, and in most preferred embodiments, at least about 95% pure. Carriers or excipients will often be added.

Solubility of a polypeptide or fragment depends upon the environment and the polypeptide. Many parameters

15

electrolyte environment, size and molecular characteristics of the polypeptide, and nature of the solvent. Typically, the temperature at which the polypeptide is used ranges from about 4° C to about 65° C. Usually the temperature at use is greater than about 18° C. For diagnostic purposes, the temperature will usually be about room temperature or warmer, but less than the denaturation temperature of components in the assay. For therapeutic purposes, the temperature will usually be body temperature, typically about 37° C for humans and mice, though under certain situations the temperature may be raised or lowered in situ or in vitro.

The size and structure of the polypeptide should generally be in a substantially stable state, and usually not in a denatured state. The polypeptide may be associated with other polypeptides in a quaternary structure, e.g., to confer solubility, or associated with lipids or detergents in a manner which approximates natural lipid bilayer interactions.

The solvent and electrolytes will usually be a biologically compatible buffer, of a type used for preservation of biological activities, and will usually approximate a physiological aqueous solvent. Usually the solvent will have a neutral pH, typically between about 5 and 10, and preferably about 7.5. On some occasions, one or more detergents will be added, typically a mild non-denaturing one, e.g., CHS (cholesteryl hemisuccinate) or CHAPS (3-[3-cholamidopropyl]dimethylammonio]-1-propane sulfonate), or a low enough concentration as to avoid significant disruption of structural or physiological properties of the protein.

III. Physical Variants

This invention also encompasses proteins or peptides having substantial amino acid sequence identity with the amino acid sequence of the 499E9. The variants include

16

Amino acid sequence homology, or sequence identity, is determined by optimizing residue matches, if necessary, by introducing gaps as required. See also Needleham, et al. (1970) J. Mol. Biol. 48:443-453;

5 Sankoff, et al. (1983) Chapter One in Time Warps, String Edits, and Macromolecules: The Theory and Practice of Sequence Comparison, Addison-Wesley, Reading, MA; and software packages from IntelliGenetics, Mountain View, CA; and the University of Wisconsin Genetics Computer

10 Group, Madison, WI. Sequence identity changes when considering conservative substitutions as matches. Conservative substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid,

15 glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine. Homologous amino acid sequences are typically intended to include natural polymorphic or allelic and interspecies variations in each respective protein sequence. Typical

20 homologous proteins or peptides will have from 25-100% identity (if gaps can be introduced), to 50-100% identity (if conservative substitutions are included) with the amino acid sequence of the 499E9. Identity measures will be at least about 35%, generally at least about 40%,

25 often at least about 50%, typically at least about 60%, usually at least about 70%, preferably at least about 80%, and more preferably at least about 90%.

The isolated 499E9 DNA can be readily modified by nucleotide substitutions, nucleotide deletions,

30 nucleotide insertions, and inversions of nucleotide stretches. These modifications result in novel DNA sequences which encode these antigens, their derivatives, or proteins having similar physiological, immunogenic, antigenic, or other functional activity. These modified

35 sequences can be used to produce mutant antigens or to enhance expression. Enhanced expression may involve gene

translation, and other mechanisms. "Mutant 499E9" encompasses a polypeptide otherwise falling within the sequence identity definition of the 499E9 as set forth above, but having an amino acid sequence which differs from that of 499E9 as normally found in nature, whether by way of deletion, substitution, or insertion. This generally includes proteins having significant identity with a protein having sequence of SEQ ID NO: 2, and as sharing various biological activities, e.g., antigenic or immunogenic, with those sequences, and in preferred embodiments contain most of the full length disclosed sequences. Full length sequences will typically be preferred, though truncated versions, e.g., soluble constructs and intact domains, will also be useful, likewise, genes or proteins found from natural sources are typically most desired. Similar concepts apply to different 499E9 proteins, particularly those found in various warm blooded animals, e.g., mammals and birds. These descriptions are generally meant to encompass all 499E9 proteins, not limited to the particular mouse embodiments specifically discussed.

499E9 mutagenesis can also be conducted by making amino acid insertions or deletions. Substitutions, deletions, insertions, or any combinations may be generated to arrive at a final construct. Insertions include amino- or carboxy- terminal fusions. Random mutagenesis can be conducted at a target codon and the expressed mutants can then be screened for the desired activity. Methods for making substitution mutations at predetermined sites in DNA having a known sequence are well known in the art, e.g., by M13 primer mutagenesis or polymerase chain reaction (PCR) techniques. See, e.g., Sambrook, et al. (1989); Ausubel, et al. (1987 and Supplements); and Kunkel, et al. (1987) Methods in Enzymol. 154:367-382.

The present invention also provides recombinant

segments from these proteins. A heterologous fusion protein is a fusion of proteins or segments which are naturally not normally fused in the same manner. A similar concept applies to heterologous nucleic acid sequences. Fusion proteins will be useful as sources for cleaving, separating, and purifying portions thereof.

In addition, new constructs may be made from combining similar functional domains from other proteins. For example, target-binding or other segments may be "swapped" between different new fusion polypeptides or fragments. See, e.g., Cunningham, et al. (1989) Science 243:1330-1336; and O'Dowd, et al. (1988) J. Biol. Chem. 263:15985-15992.

The phosphoramidite method described by Beaucage and Carruthers (1981) Tetra. Letts. 22:1859-1862, will produce suitable synthetic DNA fragments. A double stranded fragment will often be obtained either by synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence, e.g., PCR techniques.

IV. Functional Variants

The blocking of physiological response to 499E9s may result from the inhibition of binding of the antigen to its binding partner, e.g., another of itself, likely through competitive inhibition. Thus, in vitro assays of the present invention will often use isolated protein, membranes from cells expressing a membrane associated recombinant 499E9, soluble fragments comprising antigen binding segments of these proteins, or fragments attached to solid phase substrates. These assays will also allow for the diagnostic determination of the effects of either binding segment mutations and modifications, or antigen mutations and modifications, e.g., 499E9 analogs.

This invention also contemplates the use of

19

neutralizing antibodies to antigen or binding fragments compete with a test compound for binding to the protein, e.g., of natural protein sequence.

"Derivatives" of 499E9 antigens include amino acid sequence mutants from naturally occurring forms, glycosylation variants, and covalent or aggregate conjugates with other chemical moieties. Covalent derivatives can be prepared by linkage of functionalities to groups which are found in 499E9 amino acid side chains or at the N- or C- termini, e.g., by standard means. See, e.g., Lundblad and Noyes (1988) Chemical Reagents for Protein Modification, vols. 1-2, CRC Press, Inc., Boca Raton, FL; Hugli (ed.) (1989) Techniques in Protein Chemistry, Academic Press, San Diego, CA; and Wong (1991) Chemistry of Protein Conjugation and Cross Linking, CRC Press, Boca Raton, FL.

In particular, glycosylation alterations are included, e.g., made by modifying the glycosylation patterns of a polypeptide during its synthesis and processing, or in further processing steps. See, e.g., Elbein (1987) Ann. Rev. Biochem. 56:497-534. Also embraced are versions of the peptides with the same primary amino acid sequence which have other minor modifications, including phosphorylated amino acid residues, e.g., phosphotyrosine, phosphoserine, or phosphothreonine.

Fusion polypeptides between 499E9s and other homologous or heterologous proteins are also provided. Many cytokine receptors or other surface proteins are multimeric, e.g., homodimeric entities, and a repeat construct may have various advantages, including lessened susceptibility to proteolytic cleavage. Typical examples are fusions of a reporter polypeptide, e.g., luciferase, with a segment or domain of a protein, e.g., a receptor-binding segment, so that the presence or location of the fused ligand may be easily determined. See, e.g., Dull,

20

partners include bacterial β -galactosidase, trpE, Protein A, β -lactamase, alpha amylase, alcohol dehydrogenase, yeast alpha mating factor, and detection or purification tags such as a FLAG sequence or His6 sequence. See, 5 e.g., Godowski, et al. (1988) Science 241:812-816.

Fusion peptides will typically be made by either recombinant nucleic acid methods or by synthetic polypeptide methods. Techniques for nucleic acid manipulation and expression are described generally, 10 e.g., in Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual (2d ed.), vols. 1-3, Cold Spring Harbor Laboratory; and Ausubel, et al. (eds.) (1993) Current Protocols in Molecular Biology, Greene and Wiley, NY. Techniques for synthesis of polypeptides are described, 15 e.g., in Merrifield (1963) J. Amer. Chem. Soc. 85:2149-2156; Merrifield (1986) Science 232: 341-347; Atherton, et al. (1989) Solid Phase Peptide Synthesis: A Practical Approach, IRL Press, Oxford; and Grant (1992) Synthetic Peptides: A User's Guide, W.H. Freeman, NY.

20 This invention also contemplates the use of derivatives of 499E9s other than variations in amino acid sequence or glycosylation. Such derivatives may involve covalent or aggregative association with chemical moieties. Covalent or aggregative derivatives will be 25 useful as immunogens, as reagents in immunoassays, or in purification methods such as for affinity purification of binding partners, e.g., other antigens. A 499E9 can be immobilized by covalent bonding to a solid support such as cyanogen bromide-activated SEPHAROSE, by methods which 30 are well known in the art, or adsorbed onto polyolefin surfaces, with or without glutaraldehyde cross-linking, for use in the assay or purification of anti-499E9 antibodies or an alternative binding composition. The 499E9s can also be labeled with a detectable group, e.g., 35 for use in diagnostic assays. Purification of 499E9 may be effected by an immobilized antibody or complementary

A solubilized 499E9 or fragment of this invention can be used as an immunogen for the production of antisera or antibodies specific for binding to the antigen or fragments thereof. Purified antigen can be used to screen monoclonal antibodies or antigen-binding fragments, encompassing antigen binding fragments of natural antibodies, e.g., Fab, Fab', F(ab)₂, etc. Purified 499E9s can also be used as a reagent to detect antibodies generated in response to the presence of elevated levels of the antigen or cell fragments containing the antigen, both of which may be diagnostic of an abnormal or specific physiological or disease condition. This invention contemplates antibodies raised against amino acid sequences encoded by nucleotide sequence shown in SEQ ID NO: 1, or fragments of proteins containing it. In particular, this invention contemplates antibodies having binding affinity to or being raised against specific fragments which are predicted to lie outside of the lipid bilayer, both extracellular or intracellular.

The present invention contemplates the isolation of additional closely related species variants. Southern and Northern blot analysis should establish that similar genetic entities exist in other mammals. It is likely that 499E9s are widespread in species variants, e.g., rodents, lagomorphs, carnivores, artiodactyla, perissodactyla, and primates.

The invention also provides means to isolate a group of related antigens displaying both distinctness and similarities in structure, expression, and function. Elucidation of many of the physiological effects of the molecules will be greatly accelerated by the isolation and characterization of additional distinct species variants of them. In particular, the present invention provides useful probes for identifying additional homologous genetic entities in different species.

The isolated genes will allow transformation of cells lacking expression of a corresponding 499E9, e.g., either species types or cells which lack corresponding antigens and exhibit negative background activity. This should allow analysis of the function of 499E9 in comparison to untransformed control cells.

Dissection of critical structural elements which effect the various activation or differentiation functions mediated through these antigens is possible using standard techniques of modern molecular biology, particularly in comparing members of the related class. See, e.g., the homolog-scanning mutagenesis technique described in Cunningham, et al. (1989) Science 243:1339-1336; and approaches used in O'Dowd, et al. (1988) J. Biol. Chem. 263:15985-15992; and Lechleiter, et al. (1990) EMBO J. 9:4381-4390.

Intracellular functions would probably involve segments of the antigen which are normally accessible to the cytosol. However, protein internalization may occur under certain circumstances, and interaction between intracellular components and "extracellular" segments may occur. The specific segments of interaction of 499E9 with other intracellular components may be identified by mutagenesis or direct biochemical means, e.g., cross-linking or affinity methods. Structural analysis by crystallographic or other physical methods will also be applicable. Further investigation of the mechanism of signal transduction will include study of associated components which may be isolatable by affinity methods or by genetic means, e.g., complementation analysis of mutants.

Further study of the expression and control of 499E9 will be pursued. The controlling elements associated with the antigens should exhibit differential physiological, developmental, tissue specific, or other expression patterns. Upstream or downstream genetic

23

particular, physiological or developmental variants, e.g., multiple alternatively processed forms of the antigen have been found. See, e.g., SEQ ID NO: 1. Thus, differential splicing of message may lead to an

- 5 assortment of membrane bound forms, soluble forms, and modified versions of antigen.

Structural studies of the antigens will lead to design of new antigens, particularly analogs exhibiting agonist or antagonist properties on the molecule. This
10 can be combined with previously described screening methods to isolate antigens exhibiting desired spectra of activities.

V. Antibodies

- 15 Antibodies can be raised to various 499E9s, including species, polymorphic, or allelic variants, and fragments thereof, both in their naturally occurring forms and in their recombinant forms. Additionally, antibodies can be raised to 499E9s in either their active
20 forms or in their inactive forms, including native or denatured versions. Anti-idiotypic antibodies are also contemplated.

Antibodies, including binding fragments and single chain versions, against predetermined fragments of the
25 antigens can be raised by immunization of animals with conjugates of the fragments with immunogenic proteins. Monoclonal antibodies are prepared from cells secreting the desired antibody. These antibodies can be screened for binding to normal or defective 499E9s, or screened
30 for agonistic or antagonistic activity, e.g., mediated through the antigen or its binding partner. Antibodies may be agnostic or antagonistic, e.g., by sterically blocking ligand binding. These monoclonal antibodies will usually bind with at least a K_D of about 1 mM, more
35 usually at least about 300 μ M, typically at least about 100 μ M, more typically at least about 30 μ M, preferably

at least about 10 μM , and more preferably at least about 3 μM or better.

The antibodies of this invention can also be useful in diagnostic applications. As capture or non-neutralizing antibodies, they can be screened for ability to bind to the antigens without inhibiting binding by a partner. As neutralizing antibodies, they can be useful in competitive binding assays. They will also be useful in detecting or quantifying 499E9 protein or its binding partners. See, e.g., Chan (ed.) (1987) Immunology: A Practical Guide, Academic Press, Orlando, FLA; Price and Newman (eds.) (1991) Principles and Practice of Immunoassay, Stockton Press, N.Y.; and Ngo (ed.) (1988) Nonisotopic Immunoassay, Plenum Press, N.Y. Cross absorptions or other tests will identify antibodies which exhibit various spectra of specificities, e.g., unique or shared species specificities.

Further, the antibodies, including antigen binding fragments, of this invention can be potent antagonists that bind to the antigen and inhibit functional binding or inhibit the ability of a binding partner to elicit a biological response. They also can be useful as non-neutralizing antibodies and can be coupled to toxins or radionuclides so that when the antibody binds to antigen, a cell expressing it, e.g., on its surface, is killed. Further, these antibodies can be conjugated to drugs or other therapeutic agents, either directly or indirectly by means of a linker, and may effect drug targeting.

Antigen fragments may be joined to other materials, particularly polypeptides, as fused or covalently joined polypeptides to be used as immunogens. An antigen and its fragments may be fused or covalently linked to a variety of immunogens, such as keyhole limpet hemocyanin, bovine serum albumin, tetanus toxoid, etc. See Microbiology, Hoeber Medical Division, Harper and Row, 1968; Landsteiner (1962) Specificity of Serological

(1967) Methods in Immunology and Immunochemistry, vol. 1, Academic Press, New York; and Harlow and Lane (1988) Antibodies: A Laboratory Manual, CSH Press, NY, for descriptions of methods of preparing polyclonal antisera.

- 5 In some instances, it is desirable to prepare monoclonal antibodies from various mammalian hosts, such as mice, rodents, primates, humans, etc. Description of techniques for preparing such monoclonal antibodies may be found in, e.g., Stites, et al. (eds.) Basic and
- 10 Clinical Immunology (4th ed.), Lange Medical Publications, Los Altos, CA, and references cited therein; Harlow and Lane (1988) Antibodies: A Laboratory Manual, CSH Press; Goding (1986) Monoclonal Antibodies: Principles and Practice (2d ed.), Academic Press, New
- 15 York; and particularly in Kohler and Milstein (1975) in Nature 256:495-497, which discusses one method of generating monoclonal antibodies.

- Other suitable techniques involve in vitro exposure of lymphocytes to the antigenic polypeptides or
- 20 alternatively to selection of libraries of antibodies in phage or similar vectors. See, Huse, et al. (1989) "Generation of a Large Combinatorial Library of the Immunoglobulin Repertoire in Phage Lambda," Science 246:1275-1281; and Ward, et al. (1989) Nature 341:544-
- 25 546. The polypeptides and antibodies of the present invention may be used with or without modification, including chimeric or humanized antibodies. Frequently, the polypeptides and antibodies will be labeled by joining, either covalently or non-covalently, a substance
- 30 which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent
- 35 moieties, chemiluminescent moieties, magnetic particles, and the like. Patents teaching the use of such labels

26

3,996,345; 4,277,437; 4,275,149; and 4,366,241. Also, recombinant immunoglobulins may be produced, see Cabilly, U.S. Patent No. 4,816,567; Moore, et al., U.S. Patent No. 4,642,334; and Queen, et al. (1989) Proc. Nat'l Acad.

5 Sci. USA 86:10029-10033.

The antibodies of this invention can also be used for affinity chromatography in isolating the protein. Columns can be prepared where the antibodies are linked to a solid support. See, e.g., Wilchek et al. (1984)

10 Meth. Enzymol. 104:3-55.

Antibodies raised against each 499E9 will also be useful to raise anti-idiotypic antibodies. These will be useful in detecting or diagnosing various immunological conditions related to expression of the respective

15 antigens.

VI. Nucleic Acids

The described peptide sequences and the related reagents are useful in detecting, isolating, or
20 identifying a DNA clone encoding 499E9, e.g., from a natural source. Typically, it will be useful in isolating a gene from mammal, and similar procedures will be applied to isolate genes from other species, e.g., warm blooded animals, such as birds and mammals. Cross
25 hybridization will allow isolation of 499E9 from other species. A number of different approaches should be available to successfully isolate a suitable nucleic acid clone.

The purified protein or defined peptides are useful
30 for generating antibodies by standard methods, as described above. Synthetic peptides or purified protein can be presented to an immune system to generate monoclonal or polyclonal antibodies. See, e.g., Coligan (1991) Current Protocols in Immunology Wiley/Greene; and
35 Harlow and Lane (1989) Antibodies: A Laboratory Manual, Cold Spring Harbor Press. Alternatively, the 499E9 can

27

be taken of its specificity of binding, much like an antibody would be used.

For example, the specific binding composition could be used for screening of an expression library made from a cell line which expresses a 499E9. The screening can be standard staining of surface expressed antigen, or by panning. Screening of intracellular expression can also be performed by various staining or immunofluorescence procedures. The binding compositions could be used to affinity purify or sort out cells expressing the protein.

The peptide segments can also be used to predict appropriate oligonucleotides to screen a library. The genetic code can be used to select appropriate oligonucleotides useful as probes for screening. See, e.g., SEQ ID NO: 1. In combination with polymerase chain reaction (PCR) techniques, synthetic oligonucleotides will be useful in selecting correct clones from a library. Complementary sequences will also be used as probes, primers, or antisense strands. Based upon identification of the likely extracellular domain, various fragments should be particularly useful, e.g., coupled with anchored vector or poly-A complementary PCR techniques or with complementary DNA of other peptides.

This invention contemplates use of isolated DNA or fragments to encode a biologically active corresponding 499E9 polypeptide. In addition, this invention covers isolated or recombinant DNA which encodes a biologically active protein or polypeptide which is capable of hybridizing under appropriate conditions with the DNA sequences described herein. Said biologically active protein or polypeptide can be an intact antigen, or fragment, and have an amino acid sequence disclosed in, e.g., SEQ ID NO: 1. Further, this invention covers the use of isolated or recombinant DNA, or fragments thereof, which encode proteins which are homologous to a 499E9 or which was isolated using cDNA encoding a 499E9 as a

regulatory sequences in the 5' and 3' flanks, e.g., promoters, enhancers, poly-A addition signals, and others.

An "isolated" nucleic acid is a nucleic acid, e.g.,
5 an RNA, DNA, or a mixed polymer, which is substantially separated from other components which naturally accompany a native sequence, e.g., ribosomes, polymerases, and/or flanking genomic sequences from the originating species. The term embraces a nucleic acid sequence which has been
10 removed from its naturally occurring environment, and includes recombinant or cloned DNA isolates and chemically synthesized analogs or analogs biologically synthesized by heterologous systems. A substantially pure molecule includes isolated forms of the molecule.
15 Generally, the nucleic acid will be in a vector or fragment less than about 50 kb, usually less than about 30 kb, typically less than about 10 kb, and preferably less than about 6 kb.

An isolated nucleic acid will generally be a
20 homogeneous composition of molecules, but will, in some embodiments, contain minor heterogeneity. This heterogeneity is typically found at the polymer ends or portions not critical to a desired biological function or activity.

25 A "recombinant" nucleic acid is defined either by its method of production or its structure. In reference to its method of production, e.g., a product made by a process, the process is use of recombinant nucleic acid techniques, e.g., involving human intervention in the
30 nucleotide sequence, typically selection or production. Alternatively, it can be a nucleic acid made by generating a sequence comprising fusion of two fragments which are not naturally contiguous to each other, but is meant to exclude products of nature, e.g., naturally
35 occurring mutants. Thus, e.g., products made by transforming cells with any unnaturally occurring vector

29

derived using any synthetic oligonucleotide process. Such is often done to replace a codon with a redundant codon encoding the same or a conservative amino acid, while typically introducing or removing a sequence
5 recognition site.

Alternatively, it is performed to join together nucleic acid segments of desired functions to generate a single genetic entity comprising a desired combination of functions not found in the commonly available natural
10 forms. Restriction enzyme recognition sites are often the target of such artificial manipulations, but other site specific targets, e.g., promoters, DNA replication sites, regulation sequences, control sequences, or other useful features may be incorporated by design. A similar
15 concept is intended for a recombinant, e.g., fusion, polypeptide. Specifically included are synthetic nucleic acids which, by genetic code redundancy, encode polypeptides similar to fragments of these antigens, and fusions of sequences from various different species
20 variants.

A significant "fragment" in a nucleic acid context is a contiguous segment of at least about 17 nucleotides, generally at least about 22 nucleotides, ordinarily at least about 29 nucleotides, more often at least about 35
25 nucleotides, typically at least about 41 nucleotides, usually at least about 47 nucleotides, preferably at least about 55 nucleotides, and in particularly preferred embodiments will be at least about 60 or more nucleotides.

30 A DNA which codes for a 499E9 protein will be particularly useful to identify genes, mRNA, and cDNA species which code for related or homologous proteins, as well as DNAs which code for homologous proteins from different species. There are likely homologs in other
35 species, including primates, rodents, and birds. Various 499E9 homologs should be homologous and are encompassed

a more distant evolutionary relationship to the antigen can readily be isolated under appropriate conditions using these sequences if they are sufficiently homologous. Primate 499E9 proteins are of particular
5 interest.

Recombinant clones derived from the genomic sequences, e.g., containing introns, will be useful for transgenic studies, including, e.g., transgenic cells and organisms, and for gene therapy. See, e.g., Goodnow
10 (1992) "Transgenic Animals" in Roitt (ed.) Encyclopedia of Immunology, Academic Press, San Diego, pp. 1502-1504; Travis (1992) Science 256:1392-1394; Kuhn, et al. (1991) Science 254:707-710; Capecchi (1989) Science 244:1288; Robertson (1987) (ed.) Teratocarcinomas and Embryonic Stem
15 Cells: A Practical Approach, IRL Press, Oxford; and Rosenberg (1992) J. Clinical Oncology 10:180-199.

Substantial homology in the nucleic acid sequence comparison context means either that the segments, or their complementary strands, when compared, are identical
20 when optimally aligned, with appropriate nucleotide insertions or deletions, in at least about 50% of the nucleotides, generally at least about 58%, ordinarily at least about 65%, often at least about 71%, typically at least about 77%, usually at least about 85%, preferably
25 at least about 95 to 98% or more, and in particular embodiments, as high as about 99% or more of the nucleotides. Alternatively, substantial homology exists when the segments will hybridize under selective
hybridization conditions, to a strand, or its complement,
30 typically using a sequence of 499E9, e.g., in SEQ ID NO: 1. Typically, selective hybridization will occur when there is at least about 55% homology over a stretch of at least about 30 nucleotides, preferably at least about 75% over a stretch of about 25 nucleotides, and most
35 preferably at least about 90% over about 20 nucleotides. See Kanehisa (1984) Nuc. Acids Res. 12:203-213. The

longer stretches, and in certain embodiments will be over a stretch of at least about 17 nucleotides, usually at least about 28 nucleotides, typically at least about 40 nucleotides, and preferably at least about 75 to 100 or more nucleotides.

Stringent conditions, in referring to homology in the hybridization context, will be stringent combined conditions of salt, temperature, organic solvents, and other parameters, typically those controlled in hybridization reactions. Stringent temperature conditions will usually include temperatures in excess of about 30° C, usually in excess of about 37° C, typically in excess of about 55° C, preferably in excess of about 70° C. Stringent salt conditions will ordinarily be less than about 1000 mM, usually less than about 400 mM, typically less than about 250 mM, preferably less than about 150 mM. However, the combination of parameters is much more important than the measure of any single parameter. See, e.g., Wetmur and Davidson (1968) J. Mol. Biol. 31:349-370. 499E9 from other mammalian species can be cloned and isolated by cross-species hybridization of closely related species. Homology may be relatively low between distantly related species, and thus hybridization of relatively closely related species is advisable. Alternatively, preparation of an antibody preparation which exhibits less species specificity may be useful in expression cloning approaches.

VII. Making 499E9; Mimetics

DNA which encodes the 499E9 or fragments thereof can be obtained by chemical synthesis, screening cDNA libraries, or screening genomic libraries prepared from a wide variety of cell lines or tissue samples. See, e.g., Okayama and Berg (1982) Mol. Cell. Biol. 2:161-170; Gubler and Hoffman (1983) Gene 25:263-269; and Glover (ed.) (1984) DNA Cloning: A Practical Approach, IRL

herein provide useful PCR primers or allow synthetic or other preparation of suitable genes encoding a 499E9; including, naturally occurring embodiments.

5 This DNA can be expressed in a wide variety of host cells for the synthesis of a full-length 499E9 or fragments which can in turn, e.g., be used to generate polyclonal or monoclonal antibodies; for binding studies; for construction and expression of modified molecules; and for structure/function studies.

10 Vectors, as used herein, comprise plasmids, viruses, bacteriophage, integratable DNA fragments, and other vehicles which enable the integration of DNA fragments into the genome of the host. See, e.g., Pouwels, et al. (1985 and Supplements) Cloning Vectors: A Laboratory
15 Manual, Elsevier, N.Y.; and Rodriguez, et al. (1988) (eds.) Vectors: A Survey of Molecular Cloning Vectors and Their Uses, Butterworth, Boston, MA.

For purposes of this invention, DNA sequences are operably linked when they are functionally related to
20 each other. For example, DNA for a presequence or secretory leader is operably linked to a polypeptide if it is expressed as a preprotein or participates in directing the polypeptide to the cell membrane or in secretion of the polypeptide. A promoter is operably
25 linked to a coding sequence if it controls the transcription of the polypeptide; a ribosome binding site is operably linked to a coding sequence if it is positioned to permit translation. Usually, operably
30 linked means contiguous and in reading frame, however, certain genetic elements such as repressor genes are not contiguously linked but still bind to operator sequences that in turn control expression. See e.g., Rodriguez, et al., Chapter 10, pp. 205-236; Balbas and Bolivar (1990) Methods in Enzymology 185:14-37; and Ausubel, et al.
35 (1993) Current Protocols in Molecular Biology, Greene and Wiley, NY

Representative examples of suitable expression vectors include pCDNA1; pCD, see Okayama, et al. (1985) Mol. Cell Biol. 5:1136-1142; pMC1neo Poly-A, see Thomas, et al. (1987) Cell 51:503-512; and a baculovirus vector such as pAC 373 or pAC 610. See, e.g., Miller (1988) Ann. Rev. Microbiol. 42:177-199.

It will often be desired to express a 499E9 polypeptide in a system which provides a specific or defined glycosylation pattern. See, e.g., Luckow and Summers (1988) Bio/Technology 6:47-55; and Kaufman (1990) Meth. Enzymol. 185:487-511.

The 499E9, or a fragment thereof, may be engineered to be phosphatidyl inositol (PI) linked to a cell membrane, but can be removed from membranes by treatment with a phosphatidyl inositol cleaving enzyme, e.g., phosphatidyl inositol phospholipase-C. This releases the antigen in a biologically active form, and allows purification by standard procedures of protein chemistry. See, e.g., Low (1989) Biochim. Biophys. Acta 988:427-454; Tse, et al. (1985) Science 230:1003-1008; and Brunner, et al. (1991) J. Cell Biol. 114:1275-1283.

Now that the 499E9 has been characterized, fragments or derivatives thereof can be prepared by conventional processes for synthesizing peptides. These include processes such as are described in Stewart and Young (1984) Solid Phase Peptide Synthesis, Pierce Chemical Co., Rockford, IL; Bodanszky and Bodanszky (1984) The Practice of Peptide Synthesis, Springer-Verlag, New York; Bodanszky (1984) The Principles of Peptide Synthesis, Springer-Verlag, New York; and Villafranca (ed.) (1991) Techniques in Protein Chemistry II, Academic Press, San Diego, Ca.

VIII. Uses

The present invention provides reagents which will find use in diagnostic applications as described

cell mediated conditions, or below in the description of kits for diagnosis.

This invention also provides reagents with significant therapeutic value. The 499E9 (naturally occurring or recombinant), fragments thereof, and antibodies thereto, along with compounds identified as having binding affinity to 499E9, should be useful in the treatment of conditions associated with abnormal physiology or development, including abnormal proliferation, e.g., cancerous conditions, or degenerative conditions. In particular, modulation of development of lymphoid cells will be achieved by appropriate therapeutic treatment using the compositions provided herein. For example, a disease or disorder associated with abnormal expression or abnormal signaling by a 499E9 should be a likely target for an agonist or antagonist of the antigen. The antigen plays a role in regulation or development of hematopoietic cells, e.g., lymphoid cells, which affect immunological responses, e.g., autoimmune disorders.

In particular, the antigen will likely provide a costimulatory signal to cell activation. Thus, the 499E9 will likely modulate T cell mediated interactions with other cell types, e.g., cells which possess a receptor therefor. These interactions would lead, in particular contexts, to modulation of cell growth, cytokine synthesis by those or other cells, or development of particular effector cells.

Moreover, the 499E9 or antagonists could redirect T cell responses, e.g., between Th1 and Th2 polarization, or with Th0 cells. Among these agonists should be various antibodies which recognize the appropriate epitopes, e.g., which mimic binding of 499E9 to its receptor. Alternatively, they may bind to epitopes which sterically can block receptor binding.

For example, 499E9, such as the naturally

35

- may also be useful, They may provide a selective and powerful way to modulate immune responses in abnormal situations, e.g., autoimmune disorders, including rheumatoid arthritis, systemic lupus erythematosus (SLE), Hashimoto's autoimmune thyroiditis, as well as acute and chronic inflammatory responses in which T cell activation, expansion, and/or immunological T cell memory play an important role. See also Samter, et al. (eds) Immunological Diseases vols. 1 and 2, Little, Brown and Co. Regulation of T cell activation, expansion, and/or cytokine release by the naturally occurring secreted form of 499E9, or an antagonist thereof, may be effected.

- In addition, certain combination compositions with other modulators of T cell signaling would be useful. Such other signaling molecules include TcR reagents, CD40, CD40L, CTLA-8, CD28, SLAM, FAS, and their respective antagonists.

- Various abnormal conditions are known in each of the cell types shown to possess 499E9 mRNA by Northern blot analysis. See Berkow (ed.) The Merck Manual of Diagnosis and Therapy, Merck & Co., Rahway, N.J.; Thorn, et al. Harrison's Principles of Internal Medicine, McGraw-Hill, N.Y.; and Weatherall, et al. (eds.) Oxford Textbook of Medicine, Oxford University Press, Oxford. Many other medical conditions and diseases involve T cells or are T cell mediated, and many of these will be responsive to treatment by an agonist or antagonist provided herein. See, e.g., Stites and Terr (eds; 1991) Basic and Clinical Immunology Appleton and Lange, Norwalk, Connecticut; and Samter, et al. (eds) Immunological Diseases Little, Brown and Co. These problems should be susceptible to prevention or treatment using compositions provided herein.

- 499E9 antibodies can be purified and then administered to a patient, veterinary or human. These antibodies can be combined for therapeutic use with

36

conventional pharmaceutically acceptable carriers or diluents, e.g., immunogenic adjuvants, along with physiologically innocuous stabilizers, excipients, or preservatives. These combinations can be sterile
5 filtered and placed into dosage forms as by lyophilization in dosage vials or storage in stabilized aqueous preparations. This invention also contemplates use of antibodies or binding fragments thereof, including forms which are not complement binding.

10 Drug screening using 499E9 or fragments thereof can be performed to identify compounds having binding affinity to or other relevant biological effects on 499E9 functions, including isolation of associated components. Subsequent biological assays can then be utilized to
15 determine if the compound has intrinsic stimulating activity or is a blocker or antagonist in that it blocks the activity of the antigen, e.g., mutein antagonists. Likewise, a compound having intrinsic stimulating activity can activate the signal pathway and is thus an
20 agonist in that it simulates the activity of 499E9. This invention further contemplates the therapeutic use of blocking antibodies to 499E9 as antagonists and of stimulatory molecules, e.g., muteins, as agonists. This approach should be particularly useful with other 499E9
25 species variants.

The quantities of reagents necessary for effective therapy will depend upon many different factors, including means of administration, target site, physiological state of the patient, and other medicants
30 administered. Thus, treatment dosages should be titrated to optimize safety and efficacy. Typically, dosages used in vitro may provide useful guidance in the amounts useful for in situ administration of these reagents. Animal testing of effective doses for treatment of
35 particular disorders will provide further predictive information of human dosage. Various considerations are

and Gilman's: The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; and Remington's Pharmaceutical Sciences, 17th ed. (1990), Mack Publishing Co., Easton, Penn. Methods for administration are discussed therein and below, e.g., for oral, intravenous, intraperitoneal, or intramuscular administration, transdermal diffusion, and others. Pharmaceutically acceptable carriers will include water, saline, buffers, and other compounds described, e.g., in the Merck Index, Merck & Co., Rahway, New Jersey. Dosage ranges would ordinarily be expected to be in amounts lower than 1 mM concentrations, typically less than about 10 μ M concentrations, usually less than about 100 nM, preferably less than about 10 pM (picomolar), and most preferably less than about 1 fM (femtomolar), with an appropriate carrier. Slow release formulations, or a slow release apparatus will often be utilized for continuous or long term administration. See, e.g., Langer (1990) Science 249:1527-1533.

499E9, fragments thereof, and antibodies to it or its fragments, antagonists, and agonists, may be administered directly to the host to be treated or, depending on the size of the compounds, it may be desirable to conjugate them to carrier proteins such as ovalbumin or serum albumin prior to their administration. Therapeutic formulations may be administered in many conventional dosage formulations. While it is possible for the active ingredient to be administered alone, it is preferable to present it as a pharmaceutical formulation. Formulations typically comprise at least one active ingredient, as defined above, together with one or more acceptable carriers thereof. Each carrier should be both pharmaceutically and physiologically acceptable in the sense of being compatible with the other ingredients and not injurious to the patient. Formulations include those suitable for oral, rectal, nasal, topical, or parenteral administration, including intramuscular, intravenous and

conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. See, e.g., Gilman, et al. (eds.) (1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; and Remington's Pharmaceutical Sciences, 17th ed. (1990), Mack Publishing Co., Easton, Penn.; Avis, et al. (eds.) (1993) Pharmaceutical Dosage Forms: Parenteral Medications, Dekker, New York; Lieberman, et al. (eds.) (1990) Pharmaceutical Dosage Forms: Tablets, Dekker, New York; and Lieberman, et al. (eds.) (1990) Pharmaceutical Dosage Forms: Disperse Systems, Dekker, New York. The therapy of this invention may be combined with or used in association with other agents, e.g., other modulators of T cell activation, e.g., CD40, CD40 ligand, CD28, CTLA-4, B7, B70, SLAM, T cell receptor signaling entities, or their respective antagonists.

Both the naturally occurring and the recombinant form of the 499E9s of this invention are particularly useful in kits and assay methods which are capable of screening compounds for binding activity to the proteins. Several methods of automating assays have been developed in recent years so as to permit screening of tens of thousands of compounds in a short period. See, e.g., Fodor, et al. (1991) Science 251:767-773, which describes means for testing of binding affinity by a plurality of defined polymers synthesized on a solid substrate. The development of suitable assays can be greatly facilitated by the availability of large amounts of purified, soluble 499E9 as provided by this invention.

Other methods can be used to determine the critical residues in the 499E9-499E9 receptor interactions. Mutational analysis can be performed, e.g., see Somoza, et al. (1993) J. Exptl. Med. 178:549-558, to determine specific residues critical in the interaction and/or function of both extracellular domains involved in the

39

provides interactions important in intracellular signaling.

For example, antagonists can normally be found once the antigen has been structurally defined, e.g., by tertiary structure data. Testing of potential interacting analogs is now possible upon the development of highly automated assay methods using a purified 499E9. In particular, new agonists and antagonists will be discovered by using screening techniques described herein. Of particular importance are compounds found to have a combined binding affinity for a spectrum of 499E9 molecules, e.g., compounds which can serve as antagonists for species variants of 499E9.

One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant DNA molecules expressing a 499E9. Cells may be isolated which express a 499E9 in isolation from other molecules. Such cells, either in viable or fixed form, can be used for standard binding partner binding assays. See also, Parce, et al. (1989) Science 246:243-247; and Owicki, et al. (1990) Proc. Nat'l Acad. Sci. USA 87:4007-4011, which describe sensitive methods to detect cellular responses.

Another technique for drug screening involves an approach which provides high throughput screening for compounds having suitable binding affinity to a 499E9 and is described in detail in Geysen, European Patent Application 84/03564, published on September 13, 1984. First, large numbers of different small peptide test compounds are synthesized on a solid substrate, e.g., plastic pins or some other appropriate surface, see Fodor, et al. (1991). Then all the pins are reacted with solubilized, unpurified or solubilized, purified 499E9, and washed. The next step involves detecting bound 499E9.

Rational drug design may also be based upon

and other effectors or analogs. Effectors may be other proteins which mediate other functions in response to binding, or other proteins which normally interact with 499E9. One means for determining which sites interact with specific other proteins is a physical structure determination, e.g., x-ray crystallography or 2 dimensional NMR techniques. These will provide guidance as to which amino acid residues form molecular contact regions. For a detailed description of protein structural determination, see, e.g., Blundell and Johnson (1976) Protein Crystallography, Academic Press, New York.

IX. Kits

This invention also contemplates use of 499E9 proteins, fragments thereof, peptides, and their fusion products in a variety of diagnostic kits and methods for detecting the presence of another 499E9 or binding partner. Typically the kit will have a compartment containing either a defined 499E9 peptide or gene segment or a reagent which recognizes one or the other, e.g., 499E9 fragments or antibodies.

A kit for determining the binding affinity of a test compound to a 499E9 would typically comprise a test compound; a labeled compound, for example a binding partner or antibody having known binding affinity for 499E9; a source of 499E9 (naturally occurring or recombinant); and a means for separating bound from free labeled compound, such as a solid phase for immobilizing the molecule. Once compounds are screened, those having suitable binding affinity to the antigen can be evaluated in suitable biological assays, as are well known in the art, to determine whether they act as agonists or antagonists to the 499E9 signaling pathway. The availability of recombinant 499E9 polypeptides also provide well defined standards for calibrating such

A preferred kit for determining the concentration of, e.g., a 499E9 in a sample would typically comprise a labeled compound, e.g., binding partner or antibody, having known binding affinity for the antigen, a source of antigen (naturally occurring or recombinant) and a means for separating the bound from free labeled compound, e.g., a solid phase for immobilizing the 499E9. Compartments containing reagents, and instructions, will normally be provided.

Antibodies, including antigen binding fragments, specific for the 499E9 or fragments are useful in diagnostic applications to detect the presence of elevated levels of 499E9 and/or its fragments. Such diagnostic assays can employ lysates, live cells, fixed cells, immunofluorescence, cell cultures, body fluids, and further can involve the detection of antigens related to the antigen in serum, or the like. Diagnostic assays may be homogeneous (without a separation step between free reagent and antigen-binding partner complex) or heterogeneous (with a separation step). Various commercial assays exist, such as radioimmunoassay (RIA), enzyme-linked immunosorbent assay (ELISA), enzyme immunoassay (EIA), enzyme-multiplied immunoassay technique (EMIT), substrate-labeled fluorescent immunoassay (SLFIA), and the like. See, e.g., Van Vunakis, et al. (1980) Meth Enzymol. 70:1-525; Harlow and Lane (1980) Antibodies: A Laboratory Manual, CSH Press, NY; and Coligan, et al. (eds.) (1993) Current Protocols in Immunology, Greene and Wiley, NY.

Anti-idiotypic antibodies may have similar use to diagnose presence of antibodies against a 499E9, as such may be diagnostic of various abnormal states. For example, overproduction of 499E9 may result in production of various immunological reactions which may be diagnostic of abnormal physiological states, particularly in proliferative cell conditions such as cancer or

42

Frequently, the reagents for diagnostic assays are supplied in kits, so as to optimize the sensitivity of the assay. For the subject invention, depending upon the nature of the assay, the protocol, and the label, either labeled or unlabeled antibody or binding partner, or labeled 499E9 is provided. This is usually in conjunction with other additives, such as buffers, stabilizers, materials necessary for signal production such as substrates for enzymes, and the like.

Preferably, the kit will also contain instructions for proper use and disposal of the contents after use. Typically the kit has compartments for each useful reagent. Desirably, the reagents are provided as a dry lyophilized powder, where the reagents may be reconstituted in an aqueous medium providing appropriate concentrations of reagents for performing the assay.

Many of the aforementioned constituents of the drug screening and the diagnostic assays may be used without modification or may be modified in a variety of ways.

For example, labeling may be achieved by covalently or non-covalently joining a moiety which directly or indirectly provides a detectable signal. In any of these assays, the binding partner, test compound, 499E9, or antibodies thereto can be labeled either directly or indirectly. Possibilities for direct labeling include label groups: radiolabels such as ^{125}I , enzymes (U.S. Pat. No. 3,645,090) such as peroxidase and alkaline phosphatase, and fluorescent labels (U.S. Pat. No. 3,940,475) capable of monitoring the change in fluorescence intensity, wavelength shift, or fluorescence polarization. Possibilities for indirect labeling include biotinylation of one constituent followed by binding to avidin coupled to one of the above label groups.

There are also numerous methods of separating the bound 499E9 from the unbound 499E9, or alternatively the bound

immobilized on various matrixes followed by washing. Suitable matrixes include plastic such as an ELISA plate, filters, and beads. See, e.g., Coligan, et al. (eds.) (1993) Current Protocols in Immunology, Vol. 1, Chapter 2, Greene and Wiley, NY. Other suitable separation techniques include, without limitation, the fluorescein antibody magnetizable particle method described in Rattle, et al. (1984) Clin. Chem. 30:1457-1461, and the double antibody magnetic particle separation as described in U.S. Pat. No. 4,659,678.

Methods for linking proteins or their fragments to the various labels have been extensively reported in the literature and do not require detailed discussion here. Many of the techniques involve the use of activated carboxyl groups either through the use of carbodiimide or active esters to form peptide bonds, the formation of thioethers by reaction of a mercapto group with an activated halogen such as chloroacetyl, or an activated olefin such as maleimide, for linkage, or the like. Fusion proteins will also find use in these applications.

Another diagnostic aspect of this invention involves use of oligonucleotide or polynucleotide sequences taken from the sequence of a 499E9. These sequences can be used as probes for detecting levels of the 499E9 message in samples from patients suspected of having an abnormal condition, e.g., cancer or developmental problem. Since the antigen is a marker for activation, it may be useful to determine the numbers of activated T cells to determine, e.g., when additional suppression may be called for. The preparation of both RNA and DNA nucleotide sequences, the labeling of the sequences, and the preferred size of the sequences has received ample description and discussion in the literature. See, e.g., Langer-Safer, et al. (1982) Proc. Nat'l. Acad. Sci. 79:4381-4385; Caskey (1987) Science 236:962-967; and (1988) Anal. Biochem. 171:1-32.

44

Diagnostic kits which also test for the qualitative or quantitative presence of other markers are also contemplated. Diagnosis or prognosis may depend on the combination of multiple indications used as markers.

- 5 Thus, kits may test for combinations of markers. See, e.g., Viallet, et al. (1989) Progress in Growth Factor Res. 1:89-97. Other kits may be used to evaluate T cell subsets.

10 X. Methods for Isolating 499E9 Specific Binding Partners

- The 499E9 protein should interact with a receptor based, e.g., upon its similarity in structure and function to other cell surface antigens exhibiting
15 similar structure and cell type specificity of expression. Methods to isolate a receptor are made available by the ability to make purified 499E9 for screening programs. Soluble or other constructs using the 499E9 sequences provided herein will allow for
20 screening or isolation of 499E9 specific receptors. Many methods exist for expression cloning, panning, affinity isolation, or other means to identify a receptor.

- Many modifications and variations of this invention
25 can be made without departing from its spirit and scope, as will be apparent to those skilled in the art. The specific embodiments described herein are offered by way of example only, and the invention is to be limited only by the terms of the appended claims, along with the full
30 scope of equivalents to which such claims are entitled.

EXAMPLES

General Methods

- 35 Some of the standard methods are described or referenced, e.g., in Maniatis et al. (1982) Molecular

- Laboratory, Cold Spring Harbor Press; Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual (2d ed.), vols. 1-3, CSH Press, NY; Ausubel, et al., Biology, Greene Publishing Associates, Brooklyn, NY; or Ausubel, et al. (1987 and Supplements) Current Protocols in Molecular Biology, Greene and Wiley, New York; Innis, et al. (eds.) (1990) PCR Protocols: A Guide to Methods and Applications, Academic Press, N.Y. Methods for protein purification include such methods as ammonium sulfate precipitation, column chromatography, electrophoresis, centrifugation, crystallization, and others. See, e.g., Ausubel, et al. (1987 and periodic supplements); Deutscher (1990) "Guide to Protein Purification" in Methods in Enzymology vol. 182, and other volumes in this series; and manufacturer's literature on use of protein purification products, e.g., Pharmacia, Piscataway, N.J., or Bio-Rad, Richmond, CA. Combination with recombinant techniques allow fusion to appropriate segments, e.g., to a FLAG sequence or an equivalent which can be fused via a protease-removable sequence. See, e.g., Hochuli (1989) Chemische Industrie 12:69-70; Hochuli (1990) "Purification of Recombinant Proteins with Metal Chelate Absorbent" in Setlow (ed.) Genetic Engineering, Principle and Methods 12:87-98, Plenum Press, N.Y.; and Crowe, et al. (1992) OIAexpress: The High Level Expression & Protein Purification System QUIAGEN, Inc., Chatsworth, CA. Cell culture techniques are described in Doyle, et al. (eds.) (1994) Cell and Tissue Culture: Laboratory Procedures, John Wiley and Sons, NY.
- FACS analyses are described in Melamed, et al. (1990) Flow Cytometry and Sorting Wiley-Liss, Inc., New York, NY; Shapiro (1988) Practical Flow Cytometry Liss, New York, NY; and Robinson, et al. (1993) Handbook of Flow Cytometry Methods Wiley-Liss, New York, NY.
- Fluorescent labeling of appropriate reagents was performed by standard methods.

46

EXAMPLE 1: Cloning of Mouse 499E9

Production of 3W Th1 or Th2 cells is described in Openshaw, et al. (1995) J. Exp. Med. 182:1357-1367.

Briefly, Th1 or Th2 populations were derived from CD4+ T cells stimulated with antigen and antigen presenting cells in the presence of IL-12 or IL-4. Cells were stimulated once each week for 3 weeks, then harvested and restimulated, e.g., with PMA and ionomycin for 4 h. See Murphy, et al. (1996) J. Exp. Med. 183:901-913.

Total RNA can be isolated, e.g., using the guanidine thiocyanate/CsCl gradient procedure as described by Chirgwin, et al. (1978) Biochem. 18:5294-5299. Poly(A)+ RNA is isolated using, e.g., the OLIGOTEX mRNA isolation kit (QIAGEN). Such RNA from these cells is used to synthesize first strand cDNA, e.g., by using NotI/Oligo-dT primer (Gibco-BRL, Gaithersburg, MD). Double-stranded cDNA is synthesized, ligated with BstXI adaptors, digested with NotI, size fractionated for > 0.5 kilobase pairs (kb) and ligated into the NotI/BstXI sites of pJFE-14, a derivative of the pCDSR α vector. See Takebe, et al. (1985) Mol. Cell Biol. 8:466-472. Electro-competent *E. coli* DH10 α cells (Gibco-BRL) are used for transformation.

Independent clones were randomly picked and screened by hybridization using a cocktail of known cytokine cDNA's. Plasmid DNA's were prepared from clones that did not hybridize to the cytokine probes. These clones were grouped by insert size and further characterized by DNA sequencing. Clones corresponding to the 499E9 were isolated.

EXAMPLE 2: Cellular Expression of Mouse 499E9

A probe specific for cDNA encoding mouse 499E9 is used to determine tissue distribution of message encoding the antigen. Standard hybridization probes may be used for Northern analysis of RNA from appropriate

47

physiological states, in various tissues, e.g., spleen, liver, thymus, lung, etc., or in various species.

Southern analysis of cDNA libraries may also provide valuable distribution information. Standard tissue blots

5 or species blots are commercially available. Similar techniques will be useful for evaluating diagnostic or medical conditions which may correlate with expression in various cell types.

PCR analysis using appropriate primers may also be
10 used. Antibody analysis, including immunohistochemistry or FACS, may be used to determine cellular or tissue distribution.

EXAMPLE 3: Purification of 499E9 Protein

15 Multiple transfected cell lines are screened for one which expresses the antigen, membrane bound or soluble forms, at a high level compared with other cells. Various cell lines are screened and selected for their favorable properties in handling. Natural 499E9 can be
20 isolated from natural sources, or by expression from a transformed cell using an appropriate expression vector. Purification of the expressed protein is achieved by standard procedures, or may be combined with engineered means for effective purification at high efficiency from
25 cell lysates or supernatants. FLAG or His₆ segments can be used for such purification features.

EXAMPLE 4: Isolation of Homologous 499E9 Genes

The 499E9 cDNA can be used as a hybridization probe
30 to screen a library from a desired source, e.g., a primate cell cDNA library. Many different species can be screened both for stringency necessary for easy hybridization, and for presence using a probe. Appropriate hybridization conditions will be used to
35 select for clones exhibiting specificity of cross hybridization.

499E9 cDNA library degenerate

isolation of appropriate clones. Alternatively, use of appropriate primers for PCR screening will yield enrichment of appropriate nucleic acid clones.

5 Similar methods are applicable to isolate either species, polymorphic, or allelic variants. Species variants are isolated using cross-species hybridization techniques based upon isolation of a full length isolate or fragment from one species as a probe.

10 Alternatively, antibodies raised against mouse 499E9 will be used to screen for cells which express cross-reactive proteins from an appropriate, e.g., cDNA library. The purified protein or defined peptides are useful for generating antibodies by standard methods, as described above. Synthetic peptides or purified protein
15 are presented to an immune system to generate monoclonal or polyclonal antibodies. See, e.g., Coligan (1991) Current Protocols in Immunology Wiley/Greene; and Harlow and Lane (1989) Antibodies: A Laboratory Manual Cold Spring Harbor Press. The resulting antibodies are used,
20 e.g., for screening, panning, or sorting.

EXAMPLE 5: Preparation of antibodies specific for 499E9

25 Synthetic peptides or purified protein are presented to an immune system to generate monoclonal or polyclonal antibodies. See, e.g., Coligan (1991) Current Protocols in Immunology Wiley/Greene; and Harlow and Lane (1989) Antibodies: A Laboratory Manual Cold Spring Harbor Press. Polyclonal serum, or
30 hybridomas may be prepared. In appropriate situations, the binding reagent is either labeled as described above, e.g., fluorescence or otherwise, or immobilized to a substrate for panning methods.

35 EXAMPLE 6: Isolation of a Receptor for 499E9

499E9 construct expression product can be used as

partner, e.g., receptor, by taking advantage of its specificity of binding, much like an antibody would be used. A 499E9 reagent is either labeled as described above, e.g., fluorescence or otherwise, or immobilized to
5 a substrate for panning methods.

The binding composition is used to screen an expression library made from a cell line which expresses a binding partner, i.e. receptor. Standard staining techniques are used to detect or sort intracellular or
10 surface expressed receptor, or surface expressing transformed cells are screened by panning. Screening of intracellular expression is performed by various staining or immunofluorescence procedures. See also McMahan, et al. (1991) EMBO J. 10:2821-2832.

15 Alternatively, 499E9 reagents are used to affinity purify or sort out cells expressing a receptor. See, e.g., Sambrook, et al. or Ausubel, et al.

Another strategy is to screen for a membrane bound receptor by panning. The cDNA containing receptor cDNA
20 is constructed as described above. The ligand can be immobilized and used to immobilize expressing cells. Immobilization may be achieved by use of appropriate antibodies which recognize, e.g., a FLAG sequence of a 499E9 fusion construct, or by use of antibodies raised
25 against the first antibodies. Recursive cycles of selection and amplification lead to enrichment of appropriate clones and eventual isolation of receptor expressing clones.

Phage expression libraries can be screened by 499E9.
30 Appropriate label techniques, e.g., anti-FLAG antibodies, will allow specific labeling of appropriate clones.

All citations herein are incorporated herein by reference to the same extent as if each individual
35 publication or patent application was specifically and individually indicated to be incorporated by reference.

Many modifications and variations of this invention can be made without departing from its spirit and scope, as will be apparent to those skilled in the art. The specific embodiments described herein are offered by way of example only, and the invention is to be limited by the terms of the appended claims, along with the full scope of equivalents to which such claims are entitled.

SEQUENCE SUBMISSION

10

SEQ ID NO: 1 is mouse 499E9 nucleic acid sequence.

SEQ ID NO: 2 is mouse 499E9 amino acid sequence.

15

(1) GENERAL INFORMATION:

(i) APPLICANT: Schering Corporation

(ii) TITLE OF INVENTION: Mammalian Cell Surface Antigens; Related Reagents

20

(iii) NUMBER OF SEQUENCES: 2

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Schering-Plough Corporation

25

(B) STREET: 2000 Galloping Hill Road

(C) CITY: Kenilworth

(D) STATE: New Jersey

(E) COUNTRY: USA

(F) ZIP: 07033-0530

30

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Diskette

(B) COMPUTER: Apple Macintosh

(C) OPERATING SYSTEM: Macintosh 7.5.3

35

(D) SOFTWARE: Microsoft Word 6.0

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE: 12-DEC-1997

40

(C) CLASSIFICATION:

51

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US 60/032,846

(B) FILING DATE: 13-DEC-1996

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Thampoe, Immac J..

(B) REGISTRATION NUMBER: 36,322

(C) REFERENCE/DOCKET NUMBER: DX0686 PCT

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (908)298-5061

(B) TELEFAX: (908)298-5388

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2191 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 125..1072

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GCCAGGACCT CTGTGAACCG GTCGGGGCGG GGGCCGCCTG GCCGGGAGTC TGCTCGGCGG
60

TGGGTGGCCG AGGAAGGGAG AGAACGATCG CGGAGCAGGG CGCCCGAACT CCGGGCGCCG
120

CGCC ATG CGC CGG GCC AGC CGA GAC TAC GGC AAG TAC CTG CGC AGC TCG
169

Met Arg Arg Ala Ser Arg Asp Tyr Gly Lys Tyr Leu Arg Ser Ser
1 5 10 15

GAG GAG ATG GGC AGC GGC CCC GGC GTC CCA CAC GAG GGT CCG CTG CAC
217

Glu Glu Met Gly Ser Gly Pro Gly Val Pro His Glu Gly Pro Leu His
20 25 30

CCC GCG CCT TCT GCA CCG GCT CCG GCG CCG CCA CCC GCC GCC TCC CGC
265

Pro Ala Pro Ser Ala Pro Ala Pro Ala Pro Pro Pro Ala Ala Ser Arg
35 40 45

52

TCC ATG TTC CTG GCC CTC CTG GGG CTG GGA CTG GGC CAG GTG GTC TGC
 313
 Ser Met Phe Leu Ala Leu Leu Gly Leu Gly Leu Gly Gln Val Val Cys
 50 55 60
 5
 AGC ATC GCT CTG TTC CTG TAC TTT CGA GCG CAG ATG GAT CCT AAC AGA
 361
 Ser Ile Ala Leu Phe Leu Tyr Phe Arg Ala Gln Met Asp Pro Asn Arg
 65 70 75
 10
 ATA TCA GAA GAC AGC ACT CAC TGC TTT TAT AGA ATC CTG AGA CTC CAT
 409
 Ile Ser Glu Asp Ser Thr His Cys Phe Tyr Arg Ile Leu Arg Leu His
 80 85 90 95
 15
 GAA AAC GCA GGT TTG CAG GAC TCG ACT CTG GAG AGT GAA GAC ACA CTA
 457
 Glu Asn Ala Gly Leu Gln Asp Ser Thr Leu Glu Ser Glu Asp Thr Leu
 100 105 110
 20
 CCT GAC TCC TGC AGG AGG ATG AAA CAA GCC TTT CAG GGG GCC CTC CAG
 505
 Pro Asp Ser Cys Arg Arg Met Lys Gln Ala Phe Gln Gly Ala Val Gln
 115 120 125
 25
 AAG GAA CTG CAA CAC ATT GTG GGG CCA CAG CGC TTC TCA GGA GCT CCA
 553
 Lys Glu Leu Gln His Ile Val Gly Pro Gln Arg Phe Ser Gly Ala Pro
 130 135 140
 30
 GCT ATG ATG GAA GGC TCA TGG TTG GAT GTG GCC CAG CGA GGC AAG CCT
 601
 Ala Met Met Glu Gly Ser Trp Leu Asp Val Ala Gln Arg Gly Lys Pro
 145 150 155
 35
 GAG GCC CAG CCA TTT GCA CAC CTC ACC ATC AAT GCT GCC AGC ATC CCA
 649
 Glu Ala Gln Pro Phe Ala His Leu Thr Ile Asn Ala Ala Ser Ile Pro
 160 165 170 175
 40
 TCG GGT TCC CAT AAA GTC ACT CTG TCC TCT TGG TAC CAC GAT CGA GGC
 697
 Ser Gly Ser His Lys Val Thr Leu Ser Ser Trp Tyr His Asp Arg Gly
 180 185 190
 45
 TGG GCC AAG ATC TCT AAC ATG ACG TTA AGC AAC GGA AAA CTA AGG GTT
 745
 Trp Ala Lys Ile Ser Asn Met Thr Leu Ser Asn Gly Lys Leu Arg Val
 195 200 205
 50
 AAC CAA GAT GGC TTC TAT TAC CTG TAC GCC AAC ATT TGC TTT CGG CAT
 793
 Asn Gln Asp Gly Phe Tyr Tyr Leu Tyr Ala Asn Ile Cys Phe Arg His
 210

CAT GAA ACA TCG GGA AGC GTA CCT ACA GAC TAT CTT CAG CTG ATG GTG
 841
 His Glu Thr Ser Gly Ser Val Pro Thr Asp Tyr Leu Gln Leu Met Val
 225 230 235
 5
 TAT GTC GTT AAA ACC AGC ATC AAA ATC CCA AGT TCT CAT AAC CTG ATG
 889
 Tyr Val Val Lys Thr Ser Ile Lys Ile Pro Ser Ser His Asn Leu Met
 240 245 250 255
 10
 AAA GGA GGG AGC ACG AAA AAC TGG TCG GGC AAT TCT GAA TTC CAC TTT
 937
 Lys Gly Gly Ser Thr Lys Asn Trp Ser Gly Asn Ser Glu Phe His Phe
 260 265 270
 15
 TAT TCC ATA AAT GTT GGG GGA TTT TTC AAG CTC CGA GCT GGT GAA GAA
 985
 Tyr Ser Ile Asn Val Gly Gly Phe Phe Lys Leu Arg Ala Gly Glu Glu
 275 280 285
 20
 ATT AGC ATT CAG GTG TCC AAC CCT TCC CTG CTG GAT CCG GAT CAA CAT
 1033
 Ile Ser Ile Gln Val Ser Asn Pro Ser Leu Leu Asp Pro Asp Gln Asp
 290 295 300
 25
 GCG ACG TAC TTT GGG GCT TTC AAA GTT CAG GAC ATA GAC TGAGACTCAT
 1082
 Ala Thr Tyr Phe Gly Ala Phe Lys Val Gln Asp Ile Asp
 305 310 315
 30
 TTCGTGGAAC ATTAGCATGG ATGTCCTAGA TGTTTGAAAA CTTCTTAAAA AATGGATGAT
 1142
 35
 GTCTATACAT GTGTAAGACT ACTAAGAGAC ATGGCCCACG GTGTATGAAA CTCACAGCCC
 1202
 TCTCTCTTGA GCCTGTACAG GTTGTGTATA TGTAAGTCC ATAGGTGATG TTAGATTCAT
 1262
 40
 GGTGATTACA CAACGGTTTT ACAATTTTGT AATGATTTCC TAAGAATTGA ACCAGATTGG
 1322
 GAGAGGTATT CCGATGCTTA TGAAAAACTT ACACGTGAGC TATGGAAGGG GGTACACGTC
 1382
 45
 TCTGGGTCTA ACCCCTGGAC ATGTGCCACT GAGAACCCTG AAATTAAGAA GATGCCATGT
 1442
 50
 CATTGCAAAG AAATGATAGT GTGAAGGGTT AAGTTCTTTT GAATTGTTAC ATTGCGCTGG
 1502
 GACCTGCAAA TAAGTTCTTT TTTTCTAATG AGGAGAGAAA AATATATGTA TTTTATATA
 1562

54

ATGTCTAAAG TTATATTTCA GGTGTAATGT TTTCTGTGCA AAGTTTGTGA AATTATATTT
 1622

5 GTGCTATAGT ATTTGATTCA AAATAATTAA AAATGTCTCA CTCTTGACAT ATTTAATGTT
 1682

TTAAATGTAC AGATGTATTT AACTGGTGCA CTTTGTAATT CCCCTGAAGG TACTCGTAGC
 1742

10 TAAGGGGGCA GAATACTGTT TCTGGTGACC ACATGTAGTT TATTTCTTTA TTCTTTTAA
 1802

CTTAATAGAG TCTTCAGACT TGTCAAACT ATGCAAGCAA AATAAATAAA TAAAAATAAA
 1862

15 ATGAATATCT TGAATAATAA GTAGGATGTT GGTCACCAGG TGCCTTTCAA ATTTAGAAGC
 1922

TAATTGACTT TAGGAGCTGA CATAGCCAAA AAGGATACAT AATAGGCTAC TGAAAATCTG
 1982

TCAGGAGTAT TTATGCAATT ATTGAACAGG TGTCTTTTTT TACAAGAGCT ACAAATTGTA
 2042

25 AATTTTGTTC CTTTTTTTTC CCATAGAAAA TGTACTATAG TTTATCAGCC AAAAAACAAT
 2102

CCACTTTTTA ATTTAGTGAA AGTTATTTTA TTATACTGTA CAATAAAAGC ATTGTTTCTG
 2162

30 AATGGCATT TTTGGTACTT AAAAATGGC
 2191

35 (2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 316 amino acids
 (B) TYPE: amino acid
 40 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Arg Arg Ala Ser Arg Asp Tyr Gly Lys Tyr Leu Arg Ser Ser Glu
 1 5 10 15

Glu Met Gly Ser Gly Pro Gly Val Pro His Glu Gly Pro Leu His Pro
 50 20 25 30

Ala Pro Ser Ala Pro Ala Pro Ala Pro Pro Ala Ala Ser Arg Ser
 35 40 45

55

	50		55		60
	Ile Ala Leu Phe Leu Tyr Phe Arg Ala Gln Met Asp Pro Asn Arg Ile				
	65		70		75 80
5	Ser Glu Asp Ser Thr His Cys Phe Tyr Arg Ile Leu Arg Leu His Glu				
		85		90	95
	Asn Ala Gly Leu Gln Asp Ser Thr Leu Glu Ser Glu Asp Thr Leu Pro				
10		100		105	110
	Asp Ser Cys Arg Arg Met Lys Gln Ala Phe Gln Gly Ala Val Gln Lys				
		115		120	125
15	Glu Leu Gln His Ile Val Gly Pro Gln Arg Phe Ser Gly Ala Pro Ala				
		130		135	140
	Met Met Glu Gly Ser Trp Leu Asp Val Ala Gln Arg Gly Lys Pro Glu				
	145		150		155 160
20	Ala Gln Pro Phe Ala His Leu Thr Ile Asn Ala Ala Ser Ile Pro Ser				
		165		170	175
	Gly Ser His Lys Val Thr Leu Ser Ser Trp Tyr His Asp Arg Gly Trp				
25		180		185	190
	Ala Lys Ile Ser Asn Met Thr Leu Ser Asn Gly Lys Leu Arg Val Asn				
		195		200	205
30	Gln Asp Gly Phe Tyr Tyr Leu Tyr Ala Asn Ile Cys Phe Arg His His				
		210		215	220
	Glu Thr Ser Gly Ser Val Pro Thr Asp Tyr Leu Gln Leu Met Val Tyr				
	225		230		235 240
35	Val Val Lys Thr Ser Ile Lys Ile Pro Ser Ser His Asn Leu Met Lys				
		245		250	255
	Gly Gly Ser Thr Lys Asn Trp Ser Gly Asn Ser Glu Phe His Phe Tyr				
40		260		265	270
	Ser Ile Asn Val Gly Gly Phe Phe Lys Leu Arg Ala Gly Glu Glu Ile				
		275		280	285
45	Ser Ile Gln Val Ser Asn Pro Ser Leu Leu Asp Pro Asp Gln Asp Ala				
		290		295	300
	Thr Tyr Phe Gly Ala Phe Lys Val Gln Asp Ile Asp				
50		305		310	315

56

WHAT IS CLAIMED IS:

1. A substantially pure or recombinant polypeptide exhibiting at least about 85% sequence identity over a length of at least about 12 amino acids to SEQ ID NO: 2.
2. A fusion protein comprising a polypeptide of claim 1.
3. An isolated nucleic acid which encodes a polypeptide of claim 1 or fusion protein of claim 2.
4. An isolated nucleic acid having the nucleotide sequence of SEQ ID NO: 1.
5. A nucleic acid which:
 - a) hybridizes under wash conditions of 30° C and less than 2M salt to SEQ ID NO: 1; or
 - b) exhibits at least about 85% identity over a stretch of at least about 30 nucleotides to a nucleic acid encoding a 499E9 polypeptide.
6. A recombinant vector comprising a nucleic acid of either of claims 4 or 5.
7. A host cell comprising a nucleic acid or vector of any of claims 4-6.
8. A method for making a polypeptide or fusion protein comprising culturing a host cell of claim 7 under conditions in which the nucleic acid is expressed.
9. A binding compound comprising an antibody or antigen binding fragment which specifically binds to a polypeptide of claim 1.

57

10. A composition comprising a polypeptide of claim 1 or fusion protein according to claim 2.

11. A kit comprising:

- 5 a) a polypeptide of claim 1 or a fusion protein
 according to claim 2;
 b) an antibody which specifically binds to a
 polypeptide of claim 1; or
 c) a nucleic acid according to either claim 4 or 5.

10